

(4) Testing of the drug product in the same container-closure system as that in which the drug product is marketed;

(5) Testing of drug products for reconstitution at the time of dispensing (as directed in the labeling) as well as after they are reconstituted.

(b) An adequate number of batches of each drug product shall be tested to determine an appropriate expiration date and a record of such data shall be maintained. Accelerated studies, combined with basic stability information on the components, drug products, and container-closure system, may be used to support tentative expiration dates provided full shelf life studies are not available and are being conducted. Where data from accelerated studies are used to project a tentative expiration date that is beyond a date supported by actual shelf life studies, there must be stability studies conducted, including drug product testing at appropriate intervals, until the tentative expiration date is verified or the appropriate expiration date determined.

(c) For homeopathic drug products, the requirements of this section are as follows:

(1) There shall be a written assessment of stability based at least on testing or examination of the drug product for compatibility of the ingredients, and based on marketing experience with the drug product to indicate that there is no degradation of the product for the normal or expected period of use.

(2) Evaluation of stability shall be based on the same container-closure system in which the drug product is being marketed.

(d) Allergenic extracts that are labeled "No U.S. Standard of Potency" are exempt from the requirements of this section.

[43 FR 45077, Sept. 29, 1978, as amended at 46 FR 56412, Nov. 17, 1981]

§211.167 Special testing requirements.

(a) For each batch of drug product purporting to be sterile and/or pyrogen-free, there shall be appropriate laboratory testing to determine conformance to such requirements. The test procedures shall be in writing and shall be followed.

(b) For each batch of ophthalmic ointment, there shall be appropriate testing to determine conformance to specifications regarding the presence of foreign particles and harsh or abrasive substances. The test procedures shall be in writing and shall be followed.

(c) For each batch of controlled-release dosage form, there shall be appropriate laboratory testing to determine conformance to the specifications for the rate of release of each active ingredient. The test procedures shall be in writing and shall be followed.

§211.170 Reserve samples.

(a) An appropriately identified reserve sample that is representative of each lot in each shipment of each active ingredient shall be retained. The reserve sample consists of at least twice the quantity necessary for all tests required to determine whether the active ingredient meets its established specifications, except for sterility and pyrogen testing. The retention time is as follows:

(1) For an active ingredient in a drug product other than those described in paragraphs (a) (2) and (3) of this section, the reserve sample shall be retained for 1 year after the expiration date of the last lot of the drug product containing the active ingredient.

(2) For an active ingredient in a radioactive drug product, except for non-radioactive reagent kits, the reserve sample shall be retained for:

(i) Three months after the expiration date of the last lot of the drug product containing the active ingredient if the expiration dating period of the drug product is 30 days or less; or

(ii) Six months after the expiration date of the last lot of the drug product containing the active ingredient if the expiration dating period of the drug product is more than 30 days.

(3) For an active ingredient in an OTC drug product that is exempt from bearing an expiration date under §211.137, the reserve sample shall be retained for 3 years after distribution of the last lot of the drug product containing the active ingredient.

(b) An appropriately identified reserve sample that is representative of each lot or batch of drug product shall be retained and stored under conditions

§211.173

consistent with product labeling. The reserve sample shall be stored in the same immediate container-closure system in which the drug product is marketed or in one that has essentially the same characteristics. The reserve sample consists of at least twice the quantity necessary to perform all the required tests, except those for sterility and pyrogens. Except for those for drug products described in paragraph (b)(2) of this section, reserve samples from representative sample lots or batches selected by acceptable statistical procedures shall be examined visually at least once a year for evidence of deterioration unless visual examination would affect the integrity of the reserve sample. Any evidence of reserve sample deterioration shall be investigated in accordance with §211.192. The results of the examination shall be recorded and maintained with other stability data on the drug product. Reserve samples of compressed medical gases need not be retained. The retention time is as follows:

(1) For a drug product other than those described in paragraphs (b) (2) and (3) of this section, the reserve sample shall be retained for 1 year after the expiration date of the drug product.

(2) For a radioactive drug product, except for nonradioactive reagent kits, the reserve sample shall be retained for:

(i) Three months after the expiration date of the drug product if the expiration dating period of the drug product is 30 days or less; or

(ii) Six months after the expiration date of the drug product if the expiration dating period of the drug product is more than 30 days.

(3) For an OTC drug product that is exempt for bearing an expiration date under §211.137, the reserve sample must be retained for 3 years after the lot or batch of drug product is distributed.

[48 FR 13025, Mar. 29, 1983, as amended at 60 FR 4091, Jan. 20, 1995]

§211.173 Laboratory animals.

Animals used in testing components, in-process materials, or drug products for compliance with established specifications shall be maintained and controlled in a manner that assures their

21 CFR Ch. I (4–1–11 Edition)

suitability for their intended use. They shall be identified, and adequate records shall be maintained showing the history of their use.

§211.176 Penicillin contamination.

If a reasonable possibility exists that a non-penicillin drug product has been exposed to cross-contamination with penicillin, the non-penicillin drug product shall be tested for the presence of penicillin. Such drug product shall not be marketed if detectable levels are found when tested according to procedures specified in ‘Procedures for Detecting and Measuring Penicillin Contamination in Drugs,’ which is incorporated by reference. Copies are available from the Division of Research and Testing (HFD-470), Center for Drug Evaluation and Research, Food and Drug Administration, 5100 Paint Branch Pkwy., College Park, MD 20740, or available for inspection at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, or go to: http://www.archives.gov/federal_register/code_of_federal_regulations/ibr_locations.html.

[43 FR 45077, Sept. 29, 1978, as amended at 47 FR 9396, Mar. 5, 1982; 50 FR 8996, Mar. 6, 1985; 55 FR 11577, Mar. 29, 1990; 66 FR 56035, Nov. 6, 2001; 69 FR 18803, Apr. 9, 2004]

Subpart J—Records and Reports

§211.180 General requirements.

(a) Any production, control, or distribution record that is required to be maintained in compliance with this part and is specifically associated with a batch of a drug product shall be retained for at least 1 year after the expiration date of the batch or, in the case of certain OTC drug products lacking expiration dating because they meet the criteria for exemption under §211.137, 3 years after distribution of the batch.

(b) Records shall be maintained for all components, drug product containers, closures, and labeling for at least 1 year after the expiration date or, in the case of certain OTC drug